

Correlation of Radiological and Pathological Cervical Lymph Node Involvement in Oral Cavity Cancer

Asma Saleem Khan¹, Quratulain Badar², Asif Husain Osmani³, Shoaib Hanif⁴, Kaynat Siddiqui⁵

1. Resident, Radiation Oncology Department, DZUH 2. Assistant Professor, Radiation Oncology Department, DZUH 3. Consultant, Medical Oncology Department, DZUH 4. Consultant, Radiation Oncology Department DZUH 5. Resident, Radiation Oncology Department, AKUH

Corresponding author: Dr. Asma Saleem Khan, dr.asmakhan2017@gmail.com

Abstract

Objective: This study aims to detect and correlate enlarged lymph nodes radiologically with their pathological status, specifically identifying reactive and positive nodes.

Methods: This retrospective observational study was performed at Dr. Ziauddin Hospital, Karachi, Pakistan. All patients with biopsy-proven surgically resectable oral cavity cancer from January 2021 to December 2021 were included in this study. Data were gathered, including radiological and histopathological reports. The diagnostic accuracy of the scans was calculated using final histopathology as the gold standard. Data was analysed using SPSS v.26.0.

Results: 222 patients' data were reviewed, and 89 patients satisfied the inclusion criteria. The sensitivity and specificity of the CT scan were 82.98% and 38.04%, respectively. The positive predictive value was 60%, whereas the negative predictive value was 66.67%. We found that the overall accuracy of the CT scan was 61.8%.

Conclusion: We conclude that contrast-enhanced CT scans in lower-middle-income countries remain a viable radiological approach for cervical lymph node assessment that can significantly influence decisions regarding neck dissection, impacting patient survival outcomes and improving quality of life.

Keywords: Oral cavity, Squamous cell carcinoma, Lymph node, Metastasis

Introduction

According to the GLOBOCAN 2022 data, head and Neck cancers are the second most common malignancy in Pakistan.¹ The prognosis is generally worse for advanced disease. The most significant prognostic factor is the presence of cervical lymph node metastasis.^{2,3} In oral cavity SCC, the incidence of neck metastasis ranges between 34% and 50%.⁴ The survival rate drops to 50% with the increasing risk of cervical nodal involvement.⁵

Lymph node enlargement in many cases is reactive rather than metastatic, making it challenging for the treating physician to distinguish, as it influences treatment choice and prognosis.^{6,7} Different imaging modalities can be used to evaluate cervical lymphadenopathy. Each of these offers various advantages and limitations. An ultrasound scan is a useful tool to assess a lymph node 5mm or more in size and superficial. For nodes that are 2mm or smaller in size or present in deeper lymphatic chains, Computed Tomography (CT) scan offers better visualisation and has a high sensitivity.⁸ Despite offering excellent soft-tissue detail, MRI is limited by cost and scan duration.⁹ PET-CT combines metabolic imaging with anatomical localisation, making it valuable for detecting metastatic cervical lymphadenopathy. It relies on FDG uptake to distinguish reactive from metastatic lymph nodes. While effective for assessing metastases, PET-CT can yield false positives in inflammatory cases and is less accessible due to high cost and specialised facilities.^{8,9} In the management of resectable, node-positive oral cavity cancers, the current standard treatment involves excision of the primary tumour and neck dissection, followed by adjuvant radiotherapy or concurrent chemoradiotherapy (CCRT).¹⁰ For inoperable cases, definitive CCRT remains the preferred approach.¹¹

Contributions:

ASK, QB, AHO, SH, KS - Conception, Design
ASK, QB, AHO, SH, KS - Acquisition, Analysis, Interpretation
ASK, QB, AHO, SH, KS - Drafting
ASK, QB, AHO, SH, KS - Critical Review

All authors approved the final version to be published & agreed to be accountable for all aspects of the work.

Conflicts of Interest: None

Financial Support: None to report

Potential Competing Interests: None to report

Institutional Review Board Approval

8690524ASRAD

25-06-2024

Ziauddin University

Review began 11/01/2024

Review ended 07/06/2025

Published 29/09/2025

© Copyright 2025

Khan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License CC-BY-SA 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



How to cite this article: Khan AS, Badar Q, Osmani AH, Hanif S, Siddiqui K. Correlation of Radiological and Pathological Cervical Lymph Node Involvement in Oral Cavity Cancer. JRM. 2025 Sep. 29;29(3).
<https://doi.org/10.37939/jrmc.v29i3.2821>

We aim to correlate radiologically and pathologically positive cervical lymph nodes in patients with resectable oral cavity squamous cell carcinoma (SCC) in this study.

Materials And Methods

This retrospective observational study was conducted at Dr. Ziauddin Hospital, Karachi, Pakistan. Ethical approval was obtained with a waiver from the Ethics Review Committee (ERC) under reference code 8690524ASRAD. We included data from patients with biopsy-confirmed, surgically resectable oral cavity cancer treated between January 2021 and December 2021. A flowchart illustrating the inclusion and exclusion criteria was developed to provide a clear overview of the patient selection process (Figure 1). We reviewed patient files, including radiological and histopathological reports. Radiological assessment primarily uses contrast-enhanced computed tomography (CT) scans. Lymph nodes were classified as radiologically positive (cN+) if they measured greater than 10mm in short-axis diameter, exhibited enhancement similar to the primary tumour, or showed central necrosis. Data analysis was performed using IBM SPSS version 26. Descriptive statistics were calculated for patient demographics and tumour characteristics. To examine the correlation between radiologically and pathologically positive lymph nodes, we used the Pearson chi-square test. Statistical significance was defined as a p-value of <0.05.




Figure 1: Flow chart of patients who met the inclusion and exclusion criteria for the study.

Results

Of the 222 patient records reviewed, 89 met the inclusion criteria. Table 1 summarises patient and tumour characteristics. Among these, 83.1% (n=74) were male, and 16.9% (n=15) were female, with a mean age of 49.2 ± 12.06 years (range: 20–76 years). Primary tumour sites included the tongue in 49.6% (n=44) of cases and the buccal mucosa in 50.4% (n=45). Radiological staging showed that 37.1% (n=33) of patients had T2 disease and 29.2% (n=26) had T3 disease. Most patients (91%, n=81) underwent unilateral neck dissection, while 9% (n=8) had bilateral neck dissection. Radiological nodal staging identified 27% (n=24) with cN0 disease, 27% (n=24) with cN1 disease, and 46.1% (n=42) with cN2 disease, with no cases of cN3 disease. The majority (53.9%, n=48) were classified as stage IVA. (Table 1). Pathological examination after neck dissection revealed that 47.2% (n=42)

of patients were node-negative (pN0), while 52.8% (n=47) were node-positive (pN+). Among the pN+ cases, 59.6% (n=28) had primary tongue cancer, and 40.4% (n=19) had primary buccal mucosa cancer (Table 2).

Table 1: Patient and tumour characteristics

		Count (n)	Column (n% %)	Mean±SD
Gender	Male	74	83.1%	49.21±12.06
	Female	15	16.9%	
Age				
Primary site	Tongue	44	49.4%	
	Buccal mucosa	45	50.6%	
Radiological T stage (cT)	T1	13	14.6%	
	T2	33	37.1%	
	T3	26	29.2%	
	T4a	17	19.1%	
	T4b	0	0.0%	
Radiological N stage (cN)	N0	24	27.0%	
	N1	24	27.0%	
	N2	41	46.1%	
	N3	0	0.0%	
Radiological Stage	Stage I	8	9.0%	
	Stage II	15	16.9%	
	Stage III	18	20.2%	
	Stage IVa	48	53.9%	
	Stage IV b	0	0.0%	
Neck Dissection	Unilateral	81	91%	
	Bilateral	8	9%	

n represents the number of patients, % represents the percentage, and ±SD represents the standard deviation. (cT) represents the radiological tumour stage, and (cN) represents the radiological nodal stage

Table 2: Correlation between primary site and pathological nodal status

			Pathological nodal status (pN)		Total
			Positive	Negative	
Primary site	Tongue	Count (n)	28	16	44
		% within Pathological Positive	59.6%	38.1%	49.4%
	Buccal mucosa	Count (n)	19	26	45
		% within Pathological Positive	40.4%	61.9%	50.6%
Total	Count (n)		47	42	89
	% within Pathological Positive		100.0%	100.0%	100.0%

pN represents the pathological nodal status, *n* represents the number, and % denotes the percentage of the patients.

Table 3: Statistically significant correlation between radiological and histopathological lymph nodal status (p = 0.02)

			Pathological nodal status (pN)		Total	p-value*
			Positive	Negative		
Radiological nodal status (cN)	Positive	Count (n)	39	26	65	0.02
		% within Radiological positive	60.0%	40.0%	100.0%	
	Negative	Count (n)	8	16	24	
		% within Radiological positive	33.3%	66.7%	100.0%	
Total	Count		47	42	89	
	% within Radiological positive		52.8%	47.2%	100.0%	

n is the number of patients and % percentage.

*p-value is significant <0.05 by Pearson Chi-Square test

Notably, 33.3% (n=8) of patients initially labelled as radiologically node-negative (cN0) were found to be node-positive (pN+) on histopathological examination (Table 3), including 16% (n=4) identified as N3 with extranodal extension (ENE+). A

significant correlation was observed between radiological and pathological nodal statuses ($p = 0.02$). The correlation between specific radiological and corresponding pathological nodal stages was even stronger, with a highly significant p -value of 0.002. (Table 4).

Table 4: Statistically significant correlation between radiological and histopathological lymph nodal stage ($p = 0.002$)

			Pathological N stage (pN)				Total	P-value*
			N0	N1	N2	N3 with ENE		
clinical N stage (cN)	N0	Count (n)	16	1	3	4	24	p=0.002
		% within pathological N stage	38.1%	7.7%	33.3%	16.0%	27.0%	
	N1	Count (n)	16	4	2	2	24	
		% within pathological N stage	38.1%	30.8%	22.2%	8.0%	27.0%	
	N2	Count (n)	10	8	4	19	41	
		% within pathological N stage	23.8%	61.5%	44.4%	76.0%	46.1%	
Total	Count (n)	42	13	9	25	89		
	% within pathological N stage	100.0%	100.0%	100.0%	100.0%	100.0%		

$p=0.002$

cN is the radiological nodal stage, pN is the pathological nodal stage, n denotes the number of patients, and % denotes the percentage of the cases. * p -value significant <0.05 by Pearson Chi-Square test

The diagnostic performance of the CT scan in detecting lymph node metastasis was evaluated:

- Sensitivity: 82.98%
- Specificity: 38.04%
- Positive Predictive Value (PPV): 60%
- Negative Predictive Value (NPV): 66.67%
- Overall Accuracy: 61.8%

These results highlight the utility and limitations of CT scans in assessing nodal metastasis in patients with oral cavity cancer.

Discussion

This study evaluated the correlation between radiological and histopathological findings in cervical lymph node assessment among patients with oral cavity squamous cell carcinoma (OCSCC). Our key findings highlight that the CT scan demonstrated a sensitivity of 82.98% and a specificity of 38.04%, with an overall diagnostic accuracy of 61.8%. Notably, 33.3% of lymph nodes that appeared radiologically negative were confirmed positive on histopathology.

These results align with international literature reporting variable diagnostic accuracy of CT scans for cervical lymph node metastasis, ranging from 38% to 96%.¹²⁻¹⁵ A recent study showed comparable CT sensitivity (83%) but a higher specificity (61.7%) and overall accuracy (73%), suggesting that while CT is reliable for detecting positive nodes, it may still miss subclinical metastases (16). Similarly, a local study found that 23% of radiologically negative nodes were positive histologically, leading to a drop in specificity.¹⁶

The prognostic impact of cervical lymph node involvement is well established; it reduces survival by up to 50%.^{17,18} Therefore, accurate nodal staging is essential for treatment planning. While clinical examination is an important initial step, it is subjective and operator dependent. Imaging, particularly contrast-enhanced CT, provides objective staging and evaluation, although limitations remain.

Our findings reinforce the importance of elective neck dissection even in radiologically negative necks, especially in early-stage OCSCC, to improve both disease-free and overall survival.¹⁹ Given that one-third of radiologically negative nodes were positive histopathologically, relying solely on imaging may lead to under-treatment.

Limitations of the study include its single-centre nature, limited sample size, and the potential for inter-observer variability in radiological interpretation. Additionally, the absence of advanced imaging modalities like PET-CT or MRI may limit generalizability.

Conclusions

In lower-middle-income countries, where financial constraints make advanced tools unapproachable, contrast-enhanced CT scans remain a viable and convenient option for assessing cervical lymph nodes. Although their specificity is low, these scans can significantly influence neck dissection decisions, improving patient survival and quality of life.

References

1. Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2024;74(3):229-63. <https://doi.org/10.3322/caac.21834>
2. Hoda N, Bc R, Ghosh S, Ks S, B VD, Nathani J. Cervical lymph node metastasis in squamous cell carcinoma of the buccal mucosa: a retrospective study on pattern of involvement and clinical analysis. *Med Oral Patol Oral Cir Bucal.* 2021;26(1):e84-e9. <https://doi.org/10.4317/medoral.24016>
3. Cao LM, Zhong NN, Li ZZ, Huo FY, Xiao Y, Liu B, et al. Lymph node metastasis in oral squamous cell carcinoma: Where we are and where we are going. *Clinical and Translational Discovery.* 2023;3(4). <https://doi.org/10.1002/ctd2.227>
4. Gogna S, Kashyap S, Gupta N. Neck cancer resection and dissection. *StatPearls.* 2023 Aug 8. https://www.statpearls.com/point-of-care/22593#ref_25889704
5. Haidari S, Obermeier KT, Kraus M, Otto S, Probst FA, Liokatis P. Nodal Disease and Survival in Oral Cancer: Is Occult Metastasis a Burden Factor Compared to Preoperatively Nodal Positive Neck Cancers (Basel). 2022 Aug 31;14(17):4241. <https://doi.org/10.3390/cancers14174241>
6. Alqutub S, Alqutub A, Bakhshwin A, Mofti Z, Alqutub S, Alkhamesi AA, Nujoom MA, Rammal A, Merdad M, Marzouki HZ. Histopathological predictors of lymph node metastasis in oral cavity squamous cell carcinoma: a systematic review and meta-analysis. *Front Oncol.* 2024 May 14;14:1401211. <https://doi.org/10.3389/fonc.2024.1401211>
7. Arun I, Maity N, Hameed S, Jain PV, Manikantan K, Sharan R, Arun P. Lymph node characteristics and their prognostic significance in oral squamous cell carcinoma. *Head & Neck.* 2021 Feb;43(2):520-33. <https://doi.org/10.1002/hed.26499>
8. Takamura M, Nikkuni Y, Hayashi T, Katsura K, Nishiyama H, Yamazaki M, et al. Comparing the Diagnostic Accuracy of Ultrasonography, CT, MRI, and PET/CT in Cervical Lymph Node Metastasis of Oral Squamous Cell Carcinoma. *Biomedicines.* 2023;11(12):3119. <https://doi.org/10.3390/biomedicines11123119>
9. Vishwanath V, Jafarieh S, Rembielak A. The role of imaging in head and neck cancer: An overview of different imaging modalities in primary diagnosis and staging of the disease. *J Contemp Brachytherapy.* 2020;12(5):512-8. <https://doi.org/10.5114/jcb.2020.100386>
10. Thoenissen P, Heselich A, Deeg S, Al-Maawi S, Tanneberger A, Sader R, et al. Extent of Neck Dissection and Cervical Lymph Node Involvement in Oral Squamous Cell Carcinoma. *Frontiers in Oncology.* <https://doi.org/10.3389/fonc.2022.812864>
11. Hosni A, Chiu K, Huang SH, Xu W, Huang J, Bayley A, et al. Non-operative management for oral cavity carcinoma: Definitive radiation therapy as a potential alternative treatment approach. *Radiotherapy and Oncology.* 2021;154:70-5. <https://doi.org/10.1016/j.radonc.2020.08.013>
12. Narayana ML, Kumarguru B, Hameed Arafath A, Gaur U, Lakshmi P, Sravani AL. Correlation of clinical, radiological and histopathological cervical lymph node involvement in oral cancer. *International Journal of Otorhinolaryngology and Head and Neck Surgery.* 2020;6(2):311. <http://dx.doi.org/10.18203/issn.2454-5929.ijohns20200143>
13. Fu JY, Zhu L, Li J, Chen PQ, Zhang CY, Shi WT, et al. Value of preoperative computed tomography in predicting the local staging of cancer of the tongue at primary surgery. *Br J Oral Maxillofac Surg.* 2020;58(8):997-100210. <https://doi.org/10.1016/j.bjoms.2020.05.014>
14. Sharma R, Agarwal N. Comparison of CT scan and intraoperative findings of cervical lymph node metastasis in oral squamous cell carcinoma with post-operative histopathology. *Int J Otorhinolaryngol Head Neck Surg.* 2021 Jun;7:950-8. <https://doi.org/10.18203/issn.2454-5929.ijohns20212114>
15. Morisada MV, Bewley AF, Broadhead K, et al. CT predictors of sub-centimeter occult lymph node metastases in oral cavity squamous cell carcinoma: A case-control study. *The Neuroradiology Journal.* 2023;37(2):214-220. <https://doi.org/10.1177/19714009231224447>
16. Qureshi TA, Wasif M, Awan MS, Muhammad AY, Mughal A, Ameen A. Role of contrast enhanced computed tomography in assessing cervical lymph node metastases in oral cavity squamous cell carcinoma. *JPMA The Journal of the Pakistan Medical Association.* 2021;71(3):826. node, Metastasis. <https://doi.org/10.47391/JPMA.594>
17. Dolens EdS, Dourado MR, Almangush A, Salo TA, Gurgel Rocha CA, da Silva SD, et al. The impact of histopathological features on the prognosis of oral squamous cell carcinoma: a comprehensive review and meta-analysis. *Frontiers in oncology.* 2021;11:784924. <https://doi.org/10.3389/fonc.2021.784924>
18. Kim MJ, Ahn K-M. Prognostic factors of oral squamous cell carcinoma: the importance of recurrence and pTNM stage. *Maxillofacial Plastic and Reconstructive Surgery.* 2024;46(1):8. <https://doi.org/10.1186/s40902-024-00410-3>
19. Beddow T, Low TH, Gao K, Wykes J, Gupta R, Clark J, Elliott M. Elective neck dissection improves regional control in early oral squamous cell carcinoma. *Australian Journal of Otolaryngology.* 2023 Jun;6:1-9. <https://doi.org/10.21037/ajo-22-20>