

Unmasking The Culprit: Recurrent Nosebleeds From Masson's Tumor

Javeria Awan¹, Sundas Masood², Haitham Akaash³, Sadia Chaudhary⁴

Abstract

Summary: Masson's tumor, also called Intravascular papillary endothelial hyperplasia (IPEH), is a benign condition affecting various parts of the body; nonetheless, sinonasal cavity involvement is uncommon. We present the case of a 32-year-old male who experienced recurrent epistaxis and frontal headaches for 13 years. Despite multiple nasal surgeries, his symptoms persisted without a confirmed diagnosis. Physical examination revealed numerous adhesions near the middle turbinate, which were more pronounced on the left side. A contrast-enhanced CT scan revealed a heterogeneously enhancing soft-tissue lesion in the left maxillary sinus that extended into the nasal cavity, with erosion of the sinus walls. Histopathology confirmed Masson's tumor. The patient underwent several surgeries, including lateral rhinotomy and excision of a fleshy mass in the maxillary cavity. One-year post-surgery, there was no recurrence or re-bleeding. IPEH of the sinonasal region should be part of the differential diagnosis of recurrent epistaxis; otherwise, it can present an ENT surgeon with a diagnostic conundrum.

Keywords: Nasal cavity, Epistaxis, Maxillary sinus, Nasal surgical procedures, Vascular malformations.

Introduction

Masson's tumor, also known as intravascular papillary endothelial hyperplasia, is a rare entity that originates in various sites of the body, including the Head and Neck, digits, and trunk. Until now, only a few cases have been reported in the Sinonasal cavity, particularly the maxillary sinus. It is a benign vascular malformation that results in thrombus formation, inflammation, and vascular stasis within a vessel, leading to endothelial cell proliferation.¹ This paper describes a case of recurrent epistaxis diagnosed as Intravascular papillary endothelial hyperplasia of the Maxillary sinus.

Case Presentation

A 32-year-old male, known hypertensive, presented with recurrent epistaxis for the past 13 years. It was associated with a Frontal Headache. There was no history of trauma or Visual Disturbances. The patient underwent multiple Nasal surgeries over the course of 13 years, which led to no relief of his symptoms nor a confirmed diagnosis of his pathology. The physical examination revealed no definite mass in the nasal cavities; however, multiple adhesions were noted near the middle turbinate, which were more pronounced on the left side.

Investigations

A computed tomography scan with contrast showed a heterogeneously enhanced soft tissue lesion measuring about 2.7 x 2.3 x 1.8 cm in the left maxillary sinus. It is shown to expand into the left nasal cavity, accompanied by erosion of the medial and anterior walls of the maxillary sinus. The histological analysis indicated intravascular proliferation of reactive endothelial cells, resulting in many papillary forms lined by a single layer of endothelial cells, thus validating the diagnosis of Masson's tumor.

Treatment

The patient underwent multiple surgeries, including a previous lateral rhinotomy. In addition, his anterior and medial maxillary walls were deficient, providing complete access to the maxillary cavity. A visible fleshy mass was present along the anterior half of the cavity, which was then excised. A nasal endoscopy was done, and a point of origin in the apex of the maxillary cavity was identified and removed. There has been no recurrence or rebleeding at a one-year follow-up.

Discussion

Intravascular papillary endothelial hyperplasia (IPEH), also known as Masson's tumor, is an uncommon benign vascular lesion characterized by reactive endothelial proliferation and thrombus formation. Since its initial description in 1923, it has been documented in numerous anatomical sites; however, involvement of the sinonasal cavity remains exceedingly rare. Recent reviews consistently highlight the infrequency of maxillary sinus involvement, with fewer than 20 cases documented in the English literature to date.^{3,11,12} IPEH typically arises within the lumen of dilated vessels, pre-existing vascular malformations, or organizing hematomas, supporting its classification as a reactive rather than neoplastic process.² Contemporary studies reaffirm that endothelial proliferation is secondary to vascular stasis and thrombosis, rather than uncontrolled cellular atypia.^{3,11}

Contributions:

JA SM - Conception, Design
JA HA SC - Acquisition, Analysis, Interpretation
JA - Drafting
JA SM HA SC - Critical Review

All authors approved the final version to be published & agreed to be accountable for all aspects of the work.

Conflicts of Interest: None

Financial Support: None to report

Potential Competing Interests:

None to report

Institutional Review Board

Approval

4153/RTH, RWP

24-07-2025

Rawalpindi Teaching Hospital

Review began 03/10/2025

Review ended 22/01/2025

Published 31/01/2026

© Copyright 2026

Awan et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY-SA 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



How to cite this article: Awan J, Masood S, Akaash H, Chaudhary S. Unmasking The Culprit: Recurrent Nosebleeds From Masson's Tumor. JRCM. 2026 Feb. 14;1(1).

<https://doi.org/10.37939/jrcm.v1i1.3059>

Clinical Case Reports



Figure 1 Axial CT showing a soft tissue lesion in the left maxillary sinus extending into the left nasal cavity



Figure 2 (A) shows an Open lateral rhinotomy owing to past surgeries



Figure 2 (B) shows an anteriorly deficient Maxillary wall on an Open Lateral rhinotomy.




Figure 3: Histopathological image showing papillary structures in a vessel (H&E x 40)

Clinically, sinonasal IPEH presents with nonspecific symptoms depending on the site and extent of involvement. Recurrent epistaxis remains the most frequently reported presenting complaint, followed by nasal obstruction, rhinorrhea, facial pain, headache, epiphora, and anosmia.¹¹ Radiologically, IPEH poses a diagnostic challenge due to its aggressive-appearing features. Recent reports consistently describe heterogeneously enhancing sinonasal masses on contrast-enhanced CT, often accompanied by bone remodeling or erosion, findings that may raise suspicion for malignancy. MRI has been shown to provide superior soft-tissue characterization, typically demonstrating intermediate T1 and heterogeneous T2 signal intensities with contrast enhancement, aiding in surgical planning and assessment of local extension.¹²

Table 1: Reports on maxillary IPEH in literature

Author	Year of publication	Age/sex	Location	Symptoms	Imaging	Surgical techniques
Stern et al. ⁶	1991	17/M	Right maxillary sinus, ethmoid, and nasal cavity.	Frontal headache, pain in the right cheek, proptosis.	CT	Caldwell Luc excision.
Lancaster et al. ⁷	1998	67/F	Left maxillary sinus and ethmoid.	Left nasal blockage, rhinorrhea, postnasal discharge, cheek pain.	CT	Endoscopic excision.
Wang et al. ⁸	2009	42/M	Left maxillary sinus, ethmoid, frontal, and nasal cavity.	Left nasal blockage, rhinorrhea, frontal headache, epistaxis.	CT/MRI	Endoscopic excision.
D'Aguzzo et al. ⁹	2019	67/F	Right maxillary sinus, nasal cavity.	Right cheek pain, rhinorrhea, and postnasal drip.	CT/MRI	Caldwell Luc, and endoscopic excise.
Cooke et al. ¹⁰	2020	28/M	Right maxillary sinus, B/L nasal cavity, and B/L ethmoid sinus	B/L nasal blockage, epistaxis, headache, Right cheek pain, and itchy eyes.	CT	Endoscopic excision and skull base repair.
Voruz F et al. ³	2020	46/M	Left maxillary sinus	Left nasal blockage and bloody serous rhinorrhea.	CT/MRI	Endoscopic excision.
Voruz F et al. ³	2020	76/M	Right maxillary sinus, nasal cavity.	Epistaxis and rhinorrhea.	CT/MRI	Endoscopic excision.
Voruz F et al. ³	2020	33/F	Right maxillary sinus.	Rhinorrhea, orbital pressure, and headache.	CT/MRI	Endoscopic excision.
Nakamura et al. ¹¹	2023	58/M	Left maxillary sinus, B/L nasal cavity, and B/L ethmoid sinus.	B/L nasal blockage, epiphora, epistaxis.	CT/MRI	Endoscopic excision.
Maiti et al. ¹²	2025	29Y/M	Right nasal cavity, nasopharynx, maxillary sinus	Right-sided nasal obstruction, blood-tinged mucopurulence	CT/MRI	Endoscopic excision
Present case	2025	32/M	Left maxillary sinus and left nasal cavity.	Epistaxis.	CT	A combined approach of open lateral Rhinotomy and endoscopic excision.

The differential diagnosis of sinonasal IPEH is broad and includes angiosarcoma, inverted papilloma, inflammatory polyps, lymphoma, squamous cell carcinoma, and metastatic disease.⁴ Among these, angiosarcoma represents the most critical diagnostic pitfall due to its close histologic resemblance. Recent literature highlights the importance of correlating radiologic, histopathologic, and immunohistochemical findings to avoid overtreatment.^{9,11}

Definitive diagnosis relies on histopathological examination, which typically reveals papillary fronds composed of a single layer of bland endothelial cells lining fibrinous cores, with minimal atypia and absence of necrosis. Immunohistochemistry consistently demonstrates positivity for endothelial markers such as CD31, CD34, and factor VIII-related antigen, findings that have been reconfirmed in recent series.^{3,5,12}

Surgical excision remains the treatment of choice, with complete resection being curative in nearly all reported cases. Recent studies support endoscopic excision as the preferred approach when feasible, offering excellent visualization with minimal morbidity. However, open or combined approaches may be necessary in cases with extensive disease, previous surgeries, or compromised anatomy, as demonstrated in our patient.^{11,12} Recurrence is exceedingly rare and is typically associated with incomplete excision.

This case underscores the importance of considering IPEH in the differential diagnosis of long-standing recurrent epistaxis, particularly when imaging reveals a vascular sinonasal mass with bone erosion. Awareness of this rare entity among otolaryngologists, radiologists, and pathologists is essential to ensure accurate diagnosis, appropriate surgical management, and avoidance of unnecessary aggressive treatment.

Learning Outcomes

- Despite its rarity in the nasal cavity, IPEH should be considered in cases of recurrent epistaxis and included in the differential diagnosis of hemorrhagic nasal masses.
- While radiologic imaging is crucial, a definitive diagnosis requires histopathological examination.
- The treatment involves complete surgical excision.

Author Information

1. Post Graduate Trainee, ENT, Rawalpindi Teaching Hospital 2. Assistant Professor, ENT, Benazir Bhutto Hospital, Rawalpindi 3. Assistant Professor/HOD, ENT, Rawalpindi Teaching Hospital 4. Professor/HOD, ENT Benazir Bhutto Hospital, Rawalpindi.

Corresponding author: Dr. Javeria Awan, javeriaawan111@gmail.com

References

1. Masson P. Hemangioendotheliome végétant intravasculaire. *Bull. Soc. Anat. Paris.* 1923;93:517-23.
2. Hashimoto H, Daimaru Y, Enjoji M. Intravascular papillary endothelial hyperplasia: a clinicopathologic study of 91 cases. *Am J Dermatopathol.* 1983;5(6):539-46.
3. Voruz F, Arnoux G, Serex CA, de Vito C, Landis BN. Intravascular papillary endothelial hyperplasia (Masson's tumor) of the maxillary sinus. *Braz J Otorhinolaryngol.* 2022;88(1):141-5. <https://doi.org/10.1016/j.bjorl.2020.01.002>
4. Kuo TT, Sayers CP, Rosai J. Masson's "vegetant intravascular hemangioendothelioma:" a lesion often mistaken for angiosarcoma. A study of 17 cases located in the skin and soft tissues. *Cancer.* 1976 Sep;38(3):1227-36. <https://doi.org/10.1002/1097->
5. Tuna EE, Türkay B, Kurukahvecioğlu S, Ataoğlu Ö, Eryılmaz A. Sinonasal intravascular papillary endothelial hyperplasia (Masson's tumor). *ENT Case.* 2015.
6. Stern Y, Braslavsky D, Segal K, Shpitzer T, Abraham A. Intravascular papillary endothelial hyperplasia in the maxillary sinus: a benign lesion that may be mistaken for angiosarcoma. *Arch Otolaryngol Head Neck Surg.* 1991;117(10):1182-4. <https://doi.org/10.1001/archotol.1991.01870220130024>.
7. Lancaster JL, Alderson DJ, Sherman IW, Clark AH. Papillary endothelial hyperplasia (Masson's tumour) of the maxillary sinus. *J Laryngol Otol.* 1998;112(5):500-2. <https://doi.org/10.1017/s0022215100140903>.
8. Wang ZH, Hsin CH, Chen SY, Lo CY, Cheng PW. Sinonasal intravascular papillary endothelial hyperplasia successfully treated by endoscopic excision: a case report and review of the literature. *Auris Nasus Larynx.* 2009;36(3):363-6. <https://doi.org/10.1016/j.anl.2008.06.006>
9. D'Aguanno V, Ralli M, De Virgilio A, Greco A, De Vincentiis M. The role of differential diagnosis in intravascular papillary endothelial hyperplasia of the sinonasal cavity mimicking angiosarcoma: a case report. *Oncol Lett.* 2019;17(1):1253-6. <https://doi.org/10.3892/ol.2018.9717>
10. Cooke P, Goldrich D, Illoreta AM, Salama A, Shrivastava R. Intravascular papillary endothelial hyperplasia of the maxillary sinus in a patient with tricuspid atresia. *Head Neck Pathol.* 2020;14:803-7. <https://doi.org/10.1007/s12105-019-01070-w>
11. Nakamura M, Anzai T, Ishimizu E, Ashikawa K, Inoshita A, Takata Y, Matsumoto F. Intravascular papillary endothelial hyperplasia of the maxillary sinus extending into the contralateral nasal cavity. *Eur Arch Otorhinolaryngol.* 2024;281(5):2749-53. <https://doi.org/10.1007/s00405-024-08499-y>
12. Maiti M, Dutta M, Chakrabarti I, Ali H, Mazumder S, Mukherjee A, Mallick A. Sinonasal intravascular papillary endothelial hyperplasia (Masson's tumor): report of a new patient with review of literature. *European Archives of Oto-Rhino-Laryngology.* 2025 Sep 18:1-7. <https://doi.org/10.1007/s00405-025-09686-1>